



A factsheet for health professionals

Almonds

for weight management

Overweight and obesity is a major risk factor for cardiovascular disease, type 2 diabetes, some musculoskeletal conditions and some cancers.²

Rates of obesity are rising rapidly in Australia, with approximately 2 in 3 adults and 1 in 4 children reportedly overweight and obese in 2017-18.³

Gaining excess weight is mostly preventable by living a healthy lifestyle – primarily choosing healthier foods and being more physically active.⁴

Almonds are a nutrient-dense food, rich in healthy monounsaturated fats, plant-based protein and fibre, and a myriad of different vitamins and minerals including vitamin E, calcium, iron and zinc.⁵

Despite their healthy nutritional profile, people trying to lose weight often avoid nuts (including almonds) due to their high energy density. In fact, research shows that consumption of almonds in a balanced diet does not lead to weight gain and may enhance weight loss.

Recommended daily amount

The Australian Dietary Guidelines recommends **30 grams (a handful)** of nuts, including almonds, every day.¹

What your clients and patients need to know

- Almonds are high in protein and fibre, which help to increase satiety and control appetite.
- Eating a handful of almonds (30g) every day as part of a healthy balanced diet will not lead to weight gain and may help achieve weight loss.
- Almonds are a convenient and portable snack that can be eaten anywhere and any time of day.

Despite their high energy content, evidence suggests that the inclusion of almonds in an energy-restricted diet may actually enhance weight loss.



Research findings: The effect of consuming nuts (including almonds) on body weight measures

Weight management

A randomised crossover study in 2002 examined 81 male and female participants and the effect on body weight of daily almond supplementation over six months. During two sequential six-month periods, diet, body weight and habitual exercise were evaluated and almonds were provided only during the second period. Results revealed that incorporating 54g of almonds in the diet daily for six months did not, on average, lead to significant changes in body weight.⁶

Epidemiological studies looking at nuts further support these findings, revealing no association or a negative association between nut consumption and body weight.^{7,8} In addition, a 2013 meta-analysis of 33 clinical trials found that diets enriched with nuts did not increase body weight.⁹

Strengthening these findings, a 2018 updated systematic literature review looked at seven intervention studies and eight analyses around almond consumption and weight measures.¹⁰ The review concluded that consuming almonds was not associated with adverse changes in weight-related measures (body mass index, body fat, waist circumference, waist-hip ratio or lean body mass). Therefore, current evidence continues to support the position that, in the context of a healthy diet, almond consumption does not result in weight gain.

Weight loss

Despite their high energy content, evidence suggests that the inclusion of almonds in an energy-restricted diet may actually enhance weight loss. A 2003 randomised study looked at the effect of almond consumption in the context of a 24-week weight-loss program.¹¹ The participants either consumed a liquid formula-based low-calorie diet (LCD) enriched with 84g almonds per day or a liquid-based LCD supplemented with complex carbohydrates. The results found greater reductions in weight in the almond-enriched group.

A 2012 study looked at evaluating the effects of a low-calorie, almond-enriched diet compared with a low-calorie nut-free diet on body weight and cardiovascular disease risk factors over 18 months. Both groups achieved significant short-term weight reduction, which was generally maintained at 18 months. The findings concluded that incorporating limited portions of almonds (56g per day) into a behavioral weight-loss program resulted in significant weight reduction.¹²

Improvements in body fat, body mass index, waist circumference and waist-hip ratio

A 2016 randomised controlled 12-week trial of 86 healthy adults evaluated the effects of consuming almonds as part of an energy-restricted diet on body composition, compared to a nut-free energy-restricted diet.¹³ Results revealed that moderate almond consumption (15% energy from almonds) by compliant overweight and obese individuals during energy restriction resulted in greater proportional reductions of truncal and total body fat.

Similar findings were observed in a 2014 clinical trial that investigated the effects of a balanced hypocaloric almond-enriched diet (50g per day) compared to a balanced hypocaloric nut-free diet on body weight and cardiovascular disease risk factors in 108 overweight or obese women over three months.¹⁴ Results showed that both groups (almond and nut-free) experienced significant weight reduction, with the almonds group achieving greater weight-loss, reduction in body mass index, waist circumference and waist-hip ratio.



Research findings: How almonds can help with weight management

Increased satiety and appetite control

Foods that generate strong sensations of satiety can help individuals control their appetite, eat more healthily and manage their weight.¹⁵ It is well established that kilojoule-for-kilojoule not all foods deliver the same level of satiety.¹⁶ For example, in satiety studies where comparison foods were matched for energy content, there is considerable evidence that high-protein foods are more satiating than those that are high in carbohydrate and/or fat;¹⁷ and that fibre-rich foods are more satiating than low-fibre foods.¹⁸

Whole almonds have a nutritional profile consistent with satiety, being one of the tree nuts highest in plant-based protein and fibre. Research indicates that consuming almonds can have positive effects on appetite control and does not lead to significant changes in body weight.^{6,19,20,21}

A 2015 randomised crossover study found that adding 28 or 42g of almonds to the diet as a mid-morning snack was likely to increase satiety responses in a portion-dependent manner, leading to reductions in subsequent food intake so that total energy intake over the day was not increased.²²

This supported the findings of a 2013 four-week randomised controlled trial, which assessed the outcomes of consuming 43g of almonds daily with meals or as snacks.²⁰ Results showed that overall daily energy intake was reduced to compensate for energy from the almonds and did not increase the risk of weight gain among participants.

Increased resting energy expenditure

A biological mechanism that may explain the ability of nuts (including almonds) to improve body weight measures is through having a greater thermogenic effect.²³ This means consuming nuts (including almonds) increases the amount of energy and fat used by the body at rest, leading to less fat accumulation, therefore potentially curbing weight gain. This could be because nuts (including almonds) are rich in healthy unsaturated fatty acids, which are more readily oxidized (or burned) by the body than saturated fatty acids.²⁴

Decreased fat absorption

The fat in almonds is stored within its cell walls, which has shown to impact the amount of fat available for digestion.^{25,26,27} Research has shown that when whole almonds are consumed, not all cell walls are broken down when chewing, making the fat less bioaccessible and leading to incomplete fat absorption in the gut. Instead, some of the fat passes through the body and is excreted in faeces.^{25,26} This decreased absorption of fat in the body may also be a reason why almonds do not contribute to weight gain. Further, evidence suggests that we may only absorb 70% of the total calories contained in almonds.²⁸

Further, evidence suggests
that we may only absorb
70% of the total calories
contained in almonds.

References

- 1) Department of Health and Ageing. Eat for Health: Australian Dietary Guidelines. Canberra [AU]: National Health and Medical Research Council; 2013. 53p Report No: N55.
- 2) Overweight and Obesity [Internet]. Canberra: Australian Institute of Health and Welfare; c2019 [updated 2018 Jan 17; cited 2019 Jun 17]. Available from: <https://www.aihw.gov.au/reports-data/behaviours-risk-factors/overweight-obesity/overview>
- 3) Overweight and Obesity [Internet]. Canberra: Australian Bureau of Statistics; 2017-18 [updated 2019 Feb 7; cited 2019 Jun 17]. Available from: <https://www.abs.gov.au/ausstats/abs@.nsf/Lookup/by%20Subject/4364.0.55.001~2017-18~Main%20Features~Overweight%20and%20obesity~90>
- 4) Better Health Channel – Obesity [Internet]. Victoria: State Government of Victoria; c2018 [updated 2014 May; cited 2019 Jun 17]. Available from: <https://www.betterhealth.vic.gov.au/health/HealthyLiving/obesity>
- 5) Food Standards Australia New Zealand (2019). Australian Food Composition Database – Release 1. Canberra: FSANZ. Available at www.foodstandards.gov.au
- 6) Fraser, G.E., H.W. Bennett, K.B. Jaceldo-Siegl, J. Sabaté. Effect on body weight of a free 76 kilojoule (320 calorie) daily supplement of almonds for six months. *J Am Coll Nutr* [Internet]. 2002 [cited 2019 Jun 17]; 21(3):275-283. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/12074256>
- 7) Fraser GE, Sabate J, Beeson WL & Strahan TM. A possible protective effect of nut consumption on risk of coronary heart disease. The Adventist Health Study. *Arch Intern Med* [Internet] 1992 [cited 2019 Jun 17];152:1416–1424. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/1627021>
- 8) Hu FB, Stampfer MJ, Manson JE, Rimm EB, Colditz GA, Rosner BA, Speizer FE, Hennekens CH & Willett WC. Frequent nut consumption and risk of coronary heart disease in women: prospective cohort study. *Br Med J* [Internet] 1998 [cited 2019 Jun 17];317:1341–1345. Available from: <https://www.bmj.com/content/317/7169/1341>
- 9) Flores-Mateo G, Rojas-Rueda D, Basora J, Ros E, Salas-Salvado J. Nut intake and adiposity: meta-analysis of clinical trials. *Am J Clin Nutr* [Internet] 2013 [cited 2019 Jun 17];97:1346–55. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/23595878>
- 10) Neale, E, Nolan-Clark D, Tapsell LT. The effect of nut consumption on heart health: A systematic review of the literature. Sydney (AU): Nuts for Life; 2016 unpublished.
- 11) Wien MA , Sabaté JM , Iklé DN , Cole SE , Kandeel FR. Almonds vs complex carbohydrates in a weight reduction program. *Int J Obes Relat Metab Disord* [Internet] 2003 [cited 2019 Jun 17];27: 1365 – 72. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/14574348>
- 12) Foster GD , Shantz KL , Veur SSV , Oliver TL , Lent MR , Virus A , Szapary PO , Rader DJ , Zemel BS , Gilden-Tsai A. A randomized trial of the effects of an almond-enriched, hypocaloric diet in the treatment of obesity. *Am J Clin Nutr* [Internet] 2012 [cited 2019 Jun 17];96:249–54. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/22743313>
- 13) Dhillon J, Tan SY, Mattes RD. Almond consumption during energy restriction lowers truncal fat and blood pressure in compliant overweight or obese adults. *J Nutr* [Internet] 2016 [cited 2019 Jun 17];146(12):2513-2519. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27807041>
- 14) Abazarfarid Z., Salehi M., Keshavarzi S. The effect of almonds on anthropometric measurements and lipid profile in overweight and obese females in a weight reduction program: A randomized controlled clinical trial. *J Res Med Sci* [Internet] 2014 [cited 2019 Jun 17];19:457–464. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25097630>
- 15) Halford JC, Harrold JA. Satiety-enhancing products for appetite control: science and regulation of functional foods for weight management. *Proc Nutr Soc* [Internet] 2012 [cited 2019 Jun 17];71(2):350. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22401600>
- 16) Holt SH, Miller JB, Petocz P. A satiety index of common foods. *Eur J Clin Nutr* [Internet] 1995 [cited 2019 Jun 17];49(9):675–690. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7498104>
- 17) Paddon-Jones D, Westman E, Mattes RD, Wolfe RR, Astrup A, Westerterp-Plantenga M. Protein, weight management, and satiety. *Am J Clin Nutr* [Internet] 2008 [cited 2019 Jun 17];87(5):1558S–1561S. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18469287>
- 18) Howarth NC, Saltzman E, Roberts SB. Dietary fiber and weight regulation. *Nutr Rev* [Internet] 2001 [cited 2019 Jun 17];59(5):129–139. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11396693>
- 19) Hollis J, Mattes R. Effect of chronic consumption of almonds on body weight in healthy humans. *Br J Nutr* [Internet] 2007 [cited 2019 Jun 17];98(03):651–656. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17445351>
- 20) Tan S, Mattes R. Appetitive, dietary and health effects of almonds consumed with meals or as snacks: a randomized, controlled trial. *Eur J Clin Nutr* [Internet] 2013 [cited 2019 Jun 17];67(11):1205–1214. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24084509>
- 21) Sabaté J. Nut consumption and body weight. *Am J Clin Nutr* [Internet] 2013 [cited 2019 Jun 17];78(3):647S–650S. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12936960>
- 22) Hull S, Re R, Chambers L, Echaniz A, Wickham MS. A mid-morning snack of almonds generates satiety and appropriate adjustment of subsequent food intake in healthy women. *Eur J Nutr* [Internet] 2015 [cited 2019 Jun 17];54, 803–810. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25182142>
- 23) Casas-Agustench P, Lo'pez-Uriarte P, Bullo M, Ros E, Gomez-Flores A, Salas-Salvado J. Acute effects of three high-fat meals with different fat saturations on energy expenditure, substrate oxidation and satiety. *Clin Nutr* [Internet] 2009 [cited 2019 Jun 17];28(1):39–45. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19010571>
- 24) Piers LS, Walker KZ, Stoney RM, Soares MJ, O'Dea K. The influence of the type of dietary fat on postprandial fat oxidation rates: monounsaturated (olive oil) vs saturated fat (cream). *Int J Obes Relat Metab Disord* [Internet] 2002 [cited 2019 Jun 17];26:814–21. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12037652>
- 25) Ellis PR, Kendall CW, Ren Y, Parker C, Pacy JF, Waldron KW, Jenkins DJ. Role of cell walls in the bioaccessibility of lipids in almond seeds. *A J Clin Nutr* [Internet] 2004 [cited 2019 Jun 17]; 80(3): 604-13. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15321799>
- 26) Cassady BA, Hollis JH, Fulford AD, Considine RV, Mattes RD. Mastication of almonds: effects of lipid bioaccessibility, appetite, and hormone response. *A J Clin Nutr* [Internet] 2009 [cited 2019 Jun 17]; 89(3): 794-800. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19144727>
- 27) Mandalari G, Grundy MM, Grassby T, Parker ML, Cross KL, Chessa S, Bisignano C, Barreca D, Bellocchio E, Lagana G, Butterworth PJ, Faulks RM, Wilde PJ, Ellis PR, Waldron KW. The effects of processing and mastication on almond lipid bioaccessibility using novel methods of in vitro digestion modelling and micro-structural analysis. *Br J Nutr* [Internet] 2014 [cited 2019 Jun 17]; 112(9):1521-9. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25351860>
- 28) Novotny JA, Gebauer SK, Baer DJ. Discrepancy between the Atwater factor predicted and empirically measured energy values of almonds in human diets. *A J Clin Nutr* [Internet] 2012 [cited 2019 Jun 17];96(2):296-301. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22760558>

For more information visit the Health Professional Centre at health.australianalmonds.com.au